


AMENDMENTS TO THE CLAIMS

1. (Original): The compound of the general formula (1):



wherein

W, Z and one of X and Y are N and the other one of X and Y is CR⁸;

R⁸ is H, halo, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio or halo(C₁₋₄)alkyl;

R and R² are independently H, halo, C₁₋₈ alkyl, C₁₋₈ alkoxy, C₁₋₈ alkylthio, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyano or NR³R⁴, provided that at least one of R and R² is NR³R⁴;

R¹ is halo, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl(C₁₋₆)alkyl, C₁₋₈ alkoxy, C₁₋₈ alkylthio, aryl, aryloxy, arylthio, heteroaryl, heteroaryloxy, heteroarylthio, aryl(C₁₋₄)alkyl, aryl(C₁₋₄)alkoxy, heteroaryl(C₁₋₄)alkyl, heteroaryl(C₁₋₄)alkoxy, aryl(C₁₋₄)alkylthio, heteroaryl(C₁₋₄)alkylthio, morpholino, piperidino or pyrrolidino;

R³ and R⁴ are independently H, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, aryl(C₁₋₈)alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl(C₁₋₆)alkyl, heteroaryl, heteroaryl(C₁₋₈)alkyl, NR⁵R⁶, provided that not both R³ and R⁴ are H or NR⁵R⁶, or

R³ and R⁴ together form a C₃₋₇ alkylene or C₃₋₇ alkenylene chain optionally substituted with one or more C₁₋₄ alkyl or C₁₋₄ alkoxy groups, or,

together with the nitrogen atom to which they are attached, R³ and R⁴ form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine N-(C₁₋₄)alkyl (especially N-methyl) ring; and

R⁵ and R⁶ are independently H, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, aryl(C₁₋₈)alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl(C₁₋₆)alkyl, heteroaryl or heteroaryl(C₁₋₈)alkyl;

any of the foregoing alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties (other than for R⁸) being optionally substituted with halogen, cyano, C₁₋₆ alkoxy, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxy carbonyl, C₁₋₆ haloalkoxy, C₁₋₆ alkylthio, tri(C₁₋₄)alkylsilyl, C₁₋₆ alkylamino or C₁₋₆ dialkylamino,

any of the foregoing morpholine, thiomorpholine, piperidine, piperazine and pyrrolidine rings being optionally substituted with C₁₋₄ alkyl (especially methyl), and any of the foregoing aryl or heteroaryl groups or moieties being optionally substituted with one or more substituents selected from halo, hydroxy, mercapto, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkoxy, C₂₋₆ alkenyloxy, C₂₋₆

alkynyloxy, halo(C₁₋₆)alkyl, halo(C₁₋₆)alkoxy, C₁₋₆ alkylthio, halo(C₁₋₆)alkylthio, hydroxy(C₁₋₆)alkyl, C₁₋₄ alkoxy(C₁₋₆)alkyl, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR^{'''}R^{'''}, -NHCOR^{'''}, -NHCONR^{'''}R^{'''}, -CONR^{'''}R^{'''}, -SO₂R^{'''}, -OSO₂R^{'''}, -COR^{'''}, -CR^{'''}=NR^{'''} or -N=CR^{'''}R^{'''}, in which R^{'''} and R^{'''} are independently hydrogen, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy, halo(C₁₋₄)alkoxy, C₁₋₄ alkylthio, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C₁₋₄ alkyl or C₁₋₄ alkoxy.

2. (Original): A compound according to claim 1 wherein W, Z and one of X and Y are N and the other one of X and Y is CH.

3. (Previously presented) A compound according to claim 1 wherein R² is NR³R⁴.

4. (Original) A compound according to claim 3 wherein R is halo.

5. (Previously presented) A compound according to claim 1 wherein

R³ is C₁₋₈ alkyl, halo(C₁₋₈)alkyl, hydroxy(C₁₋₈)alkyl, C₁₋₄ alkoxy(C₁₋₈)alkyl, C₁₋₄ alkoxyhalo(C₁₋₈)alkyl, tri(C₁₋₄)alkylsilyl(C₁₋₆)alkyl, C₁₋₄ alkylcarbonyl(C₁₋₈)alkyl, C₁₋₄ alkylcarbonylhalo(C₁₋₈)alkyl, phenyl(C₁₋₄)alkyl, C₂₋₈ alkenyl, halo(C₂₋₈)alkenyl, C₂₋₈ alkynyl, C₃₋₈ cycloalkyl optionally substituted with chloro, fluoro or methyl, C₃₋₈ cycloalkyl(C₁₋₄)alkyl, phenylamino, piperidino or morpholino, the phenyl ring of phenylalkyl or phenylamino being optionally substituted with one, two or three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy and halo(C₁₋₄)alkoxy; and

R⁴ is H, C₁₋₄ alkyl, halo(C₁₋₄)alkyl or amino, or

R³ and R⁴ together form a C₃₋₇ alkylene or alkenylene chain optionally substituted with methyl, or,

together with the nitrogen atom to which they are attached, R³ and R⁴ form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine N-(C₁₋₄)alkyl (especially N-methyl) ring, in which the morpholine or piperazine rings are optionally substituted with methyl.

6. (Previously presented) A compound according to claim 1 wherein

R¹ is phenyl optionally substituted with from one to five halogen atoms or with from one to three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy or halo(C₁₋₄)alkoxy, pyridyl optionally substituted with from one to four halogen atoms or with from one to three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy or halo(C₁₋₄)alkoxy, 2- or 3-thienyl optionally substituted with from one to three halogen atoms or with from one to three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy or halo(C₁₋₄)alkoxy, or piperidino or morpholino both optionally substituted with one or two methyl groups.

7. (Original) A compound according to claim 6 wherein R¹ is 2,6-difluorophenyl, 2-fluoro-6-chlorophenyl, 2,5,6-trifluorophenyl, 2,4,6-trifluorophenyl, 2,6-difluoro-4-methoxyphenyl or pentafluorophenyl.

8. (Original): A compound according to claim 1 wherein

W, Z and one of X and Y are N and the other one of X and Y is CR⁸;

R⁸ is H, halo, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio or halo(C₁₋₄)alkyl;

one of R and R² (preferably R²) is NR³R⁴ and the other is halo;

R¹ is halo, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl(C₁₋₆)alkyl, C₁₋₈ alkoxy, C₁₋₈ alkylthio, aryl, aryloxy, arylthio, heteroaryl, heteroaryloxy, heteroarylthio, aryl(C₁₋₄)alkyl, aryl(C₁₋₄)alkoxy, heteroaryl(C₁₋₄)alkyl, heteroaryl(C₁₋₄)alkoxy, aryl(C₁₋₄)alkylthio, heteroaryl(C₁₋₄)alkylthio, morpholino, piperidino or pyrrolidino;

R³ and R⁴ are independently H, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, aryl(C₁₋₈)alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl(C₁₋₆)alkyl, heteroaryl, heteroaryl(C₁₋₈)alkyl, NR⁵R⁶, provided that not both R³ and R⁴ are H or NR⁵R⁶, or

R³ and R⁴ together form a C₃₋₇ alkylene or C₃₋₇ alkenylene chain optionally substituted with one or more C₁₋₄ alkyl or C₁₋₄ alkoxy groups, or, together with the nitrogen atom to which they are attached, R³ and R⁴ form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine N-(C₁₋₄)alkyl (especially N-methyl) ring; and

R⁵ and R⁶ are independently H, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, aryl(C₁₋₈)alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl(C₁₋₆)alkyl, heteroaryl or heteroaryl(C₁₋₈)alkyl;

any of the foregoing alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties (other than for R⁸) being optionally substituted with halogen, cyano, C₁₋₆ alkoxy, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ haloalkoxy, C₁₋₆ alkylthio, tri(C₁₋₄)alkylsilyl, C₁₋₆ alkylamino or C₁₋₆ dialkylamino,

any of the foregoing morpholine, thiomorpholine, piperidine, piperazine and pyrrolidine rings being optionally substituted with C₁₋₄ alkyl (especially methyl), and

any of the aryl, heteroaryl, aryloxy or heteroaryl groups being optionally substituted with one or more substituents selected from halo, hydroxy, mercapto, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkoxy, C₂₋₆ alkenyloxy, C₂₋₆ alkynyloxy, halo(C₁₋₆)alkyl, halo(C₁₋₆)alkoxy, C₁₋₆ alkylthio, halo(C₁₋₆)alkylthio, hydroxy(C₁₋₆)alkyl, C₁₋₄ alkoxy(C₁₋₆)alkyl, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR^{'''}R^{'''}, -NHCOR^{'''}, -NHCONR^{'''}R^{'''}, -CONR^{'''}R^{'''}, -SO₂R^{'''}, -OSO₂R^{'''}, -COR^{'''}, -CR^{'''}=NR^{'''} or -N=CR^{'''}R^{'''}, in which R^{'''} and R^{'''} are independently hydrogen, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄

alkoxy, halo(C₁₋₄)alkoxy, C₁₋₄ alkylthio, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl (C₁₋₄)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C₁₋₄ alkyl or C₁₋₄ alkoxy.

9. (Original): A compound according to claim 1 wherein

W, Z and one of X and Y are N and the other one of X and Y is CR⁸;

R⁸ is H, halo, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio or halo(C₁₋₄)alkyl;

one of R and R² (preferably R²) is NR³R⁴ and the other is halo;

R¹ is halo, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl(C₁₋₆)alkyl, C₁₋₈ alkoxy, C₁₋₈ alkylthio, aryl, aryloxy, arylthio, heteroaryl, heteroaryloxy, heteroarylthio, aryl(C₁₋₄)alkyl, aryl(C₁₋₄)alkoxy, heteroaryl(C₁₋₄)alkyl, heteroaryl(C₁₋₄)alkoxy, aryl(C₁₋₄)alkylthio, heteroaryl(C₁₋₄)alkylthio, morpholino, piperidino or pyrrolidino;

R³ is C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₂₋₄ alkenyl, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl or phenylamino in which the phenyl ring is optionally substituted with one, two or three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy and halo(C₁₋₄)alkoxy; and R⁴ is H, C₁₋₄ alkyl or amino, or

R³ and R⁴ together form a C₄₋₆ alkylene chain optionally substituted with C₁₋₄ alkyl or C₁₋₄ alkoxy, or, together with the nitrogen atom to which they are attached, R³ and R⁴ form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine N-(C₁₋₄)alkyl (especially N-methyl) ring;

any of the alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties (other than for R⁸) being optionally substituted with halogen, cyano, C₁₋₆ alkoxy, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ haloalkoxy, C₁₋₆ alkylthio, tri(C₁₋₄)alkylsilyl, C₁₋₆ alkylamino or C₁₋₆ dialkylamino,

any of the foregoing morpholine, thiomorpholine, piperidine, piperazine and pyrrolidine rings being optionally substituted with C₁₋₄ alkyl (especially methyl), and

any of the aryl or heteroaryl groups or moieties being optionally substituted with one or more substituents selected from halo, hydroxy, mercapto, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkoxy, C₂₋₆ alkenyloxy, C₂₋₆ alkynyloxy, halo(C₁₋₆)alkyl, halo(C₁₋₆)alkoxy, C₁₋₆ alkylthio, halo(C₁₋₆)alkylthio, hydroxy(C₁₋₆)alkyl, C₁₋₄ alkoxy(C₁₋₆)alkyl, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR^{'''}R^{'''}, -NHCOR^{'''}, -NHCONR^{'''}R^{'''}, -CONR^{'''}R^{'''}, -SO₂R^{'''}, -OSO₂R^{'''}, -COR^{'''}, -CR^{'''}=NR^{'''} or -N=CR^{'''}R^{'''}, in which R^{'''} and R^{'''} are independently hydrogen, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy, halo(C₁₋₄)alkoxy, C₁₋₄ alkylthio, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C₁₋₄ alkyl or C₁₋₄ alkoxy.

10. (Original) A compound according to claim 1 wherein

W, Z and one of X and Y are N and the other one of X and Y is CR⁸;

R⁸ is H, halo, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio or halo(C₁₋₄)alkyl;

R and R² are independently H, halo, C₁₋₈ alkyl, C₁₋₈ alkoxy, C₁₋₈ alkylthio, C₂₋₈ alkenyl, C₂₋₈ alkynyl, cyano or NR³R⁴, provided that at least one of R and R² (preferably R²) is NR³R⁴;

R¹ is optionally substituted phenyl;

R³ and R⁴ are independently H, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, aryl(C₁₋₈)alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl(C₁₋₆)alkyl, heteroaryl, heteroaryl(C₁₋₈)alkyl, NR⁵R⁶, provided that not both R³ and R⁴ are H or NR⁵R⁶, or

R³ and R⁴ together form a C₃₋₇ alkylene or C₃₋₇ alkenylene chain optionally substituted with one or more C₁₋₄ alkyl or C₁₋₄ alkoxy groups, or, together with the nitrogen atom to which they are attached, R³ and R⁴ form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine *N*-(C₁₋₄)alkyl (especially *N*-methyl) ring; and

R⁵ and R⁶ are independently H, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, aryl(C₁₋₈)alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl(C₁₋₆)alkyl, heteroaryl or heteroaryl(C₁₋₈)alkyl;

any of the alkyl, alkenyl, alkynyl or cycloalkyl groups or moieties (other than for R⁸) being optionally substituted with halogen, cyano, C₁₋₆ alkoxy, C₁₋₆ alkylcarbonyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ haloalkoxy, C₁₋₆ alkylthio, tri(C₁₋₄)alkylsilyl, C₁₋₆ alkylamino or C₁₋₆ dialkylamino,

any of the foregoing morpholine, thiomorpholine, piperidine, piperazine and pyrrolidine rings being optionally substituted with C₁₋₄ alkyl (especially methyl), and

any of the aryl or heteroaryl groups or moieties, including the phenyl group of R¹, being optionally substituted with one or more substituents selected from halo, hydroxy, mercapto, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkoxy, C₂₋₆ alkenyloxy, C₂₋₆ alkynyloxy, halo(C₁₋₆)alkyl, halo(C₁₋₆)alkoxy, C₁₋₆ alkylthio, halo(C₁₋₆)alkylthio, hydroxy(C₁₋₆)alkyl, C₁₋₄ alkoxy(C₁₋₆)alkyl, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR^{'''}R^{'''}, -NHCOR^{'''}, -NHCONR^{'''}R^{'''}, -CONR^{'''}R^{'''}, -SO₂R^{'''}, -OSO₂R^{'''}, -COR^{'''}, -CR^{'''}=NR^{'''} or -N=CR^{'''}R^{'''}, in which R^{'''} and R^{'''} are independently hydrogen, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy, halo(C₁₋₄)alkoxy, C₁₋₄ alkylthio, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C₁₋₄ alkyl or C₁₋₄ alkoxy.

11. (Original) A compound according to claim 1 wherein

W, Z and one of X and Y are N and the other one of X and Y is CR⁸;

R⁸ is H, halo, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio or halo(C₁₋₄)alkyl;

R is H, halo, C₁₋₄ alkyl), C₁₋₄ alkoxy or cyano;

R¹ is phenyl optionally substituted with from one to five halogen atoms or with from one to three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy or halo(C₁₋₄)alkoxy, pyridyl optionally substituted with from one to four halogen atoms or with from one to three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy or halo(C₁₋₄)alkoxy, 2- or 3-thienyl optionally substituted with from one to three halogen atoms or with from one to three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy or halo(C₁₋₄)alkoxy, or piperidino or morpholino both optionally substituted with one or two methyl groups;

R² is NR³R⁴;

R³ is C₁₋₈ alkyl, halo(C₁₋₈)alkyl, hydroxy(C₁₋₈)alkyl, C₁₋₄ alkoxy(C₁₋₈)alkyl, C₁₋₄ alkoxyhalo(C₁₋₈)alkyl, tri(C₁₋₄)alkylsilyl(C₁₋₆)alkyl, C₁₋₄ alkylcarbonyl(C₁₋₈)alkyl, C₁₋₄ alkylcarbonylhalo(C₁₋₈)alkyl, phenyl(C₁₋₄)alkyl, C₂₋₈ alkenyl, halo(C₂₋₈)alkenyl, C₂₋₈ alkynyl, C₃₋₈ cycloalkyl optionally substituted with chloro, fluoro or methyl, C₃₋₈ cycloalkyl(C₁₋₄)alkyl, phenylamino, piperidino or morpholino, the phenyl ring of phenylalkyl or phenylamino being optionally substituted with one, two or three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy and halo(C₁₋₄)alkoxy; and

R⁴ is H, C₁₋₄ alkyl, halo(C₁₋₄)alkyl or amino, or

R³ and R⁴ together form a C₃₋₇ alkylene or C₃₋₇ alkenylene chain optionally substituted with methyl, or,

together with the nitrogen atom to which they are attached, R³ and R⁴ form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine *N*-(C₁₋₄)alkyl (especially *N*-methyl) ring, in which the morpholine or piperazine rings are optionally substituted with methyl.

12. (Original) A compound according to claim 1 wherein

W, Z and one of X and Y are N and the other one of X and Y is CR⁸;

R⁸ is H, halo, C₁₋₄ alkyl, C₁₋₄ alkoxy or halo(C₁₋₄)alkyl;

R is halo;


R¹ is phenyl optionally substituted with from one to five halogen atoms or with from one to three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy or halo(C₁₋₄)alkoxy;

R² is NR³R⁴;


R³ is C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₂₋₄ alkenyl, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl or phenylamino in which the phenyl ring is optionally substituted with one, two or three substituents selected from halo, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy and halo(C₁₋₄)alkoxy; and

R^4 is H, C_{1-4} alkyl or amino, or R^3 and R^4 together form a C_{4-6} alkylene chain optionally substituted with methyl, or, together with the nitrogen atom to which they are attached, R^3 and R^4 form a morpholine ring.

13. (Original) A process for preparing a compound of the general formula (1) according to claim 1 wherein one of R and R^2 is chloro or fluoro and the other is NR^3R^4 and W, X, Y, Z, R^1 , R^3 and R^4 are as defined in claim 1, which comprises reacting an amine of the general formula NR^3R^4 with a compound of the general formula (6) or (13):



14. (Withdrawn) The intermediate chemicals having the general formulae (4), (5), (6) and (13):




wherein W, X, Y, Z and R^1 are as defined in claim 1 and R^7 is C_{1-4} alkyl.

15. (Original) A plant fungicidal composition comprising a fungicidally effective amount of a compound as defined in claim 1 and a suitable carrier or diluent therefor.

16. (Previously presented) A method of combating or controlling phytopathogenic fungi which comprises applying to a plant, to a seed of a plant, to the locus of the plant or seed or to soil or to any other plant growth medium, a fungicidally effective amount of a compound according to claim 1.

17. (New) A compound according to claim 1 having the formula:



wherein:

R is H, halo, C_{1-8} alkyl, C_{1-8} alkoxy, C_{1-8} alkylthio, C_{2-8} alkenyl, C_{2-8} alkynyl, cyano or NR^3R^4 ;

R³ and R⁴ are independently H, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, aryl(C₁₋₈)alkyl, heteroaryl, heteroaryl(C₁₋₈)alkyl, NR⁵R⁶, provided that not both R³ and R⁴ are H or NR⁵R⁶, or

R³ and R⁴ together form a C₃₋₇ alkylene or C₃₋₇ alkenylene chain optionally substituted with one or more C₁₋₄ alkyl or C₁₋₄ alkoxy groups, or,

together with the nitrogen atom to which they are attached, R³ and R⁴ form a morpholine, thiomorpholine, thiomorpholine S-oxide or thiomorpholine S-dioxide ring or a piperazine or piperazine N-(C₁₋₄)alkyl (especially N-methyl) ring; and

R⁵ and R⁶ are independently H, C₁₋₈ alkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, aryl, aryl(C₁₋₈)alkyl, C₃₋₈ cycloalkyl, C₃₋₈ cycloalkyl(C₁₋₆)alkyl, heteroaryl or heteroaryl(C₁₋₈)alkyl.